

DOI 10.2478/pjvs-2013-0029

Original article

# Changes in expression of calbindin 28 kDa in the small intestine of red kidney bean (*Phaseolus vulgaris*) lectin-treated suckling piglets

A. Zacharko-Siembida<sup>1</sup>, J.L. Valverde Piedra<sup>2,3</sup>, M.B. Arciszewski<sup>1</sup>

<sup>1</sup> Department of Animal Anatomy and Histology

<sup>2</sup> Sub-Department of Toxicology and Environment Protection, Faculty of Veterinary Medicine, University of Life Sciences, Akademicka 12, 20-033, Lublin, Poland

<sup>3</sup> Former affiliation: Department of Biochemistry and Animal Physiology, Sub-Department of Animal Physiology, Faculty of Veterinary Medicine, University of Life Sciences in Lublin, Poland

## Abstract

Calbindin (CB) is a calcium binding protein playing a role in calcium uptake and anti-apoptotic cellular protection. The expression of CB was immunohistochemically studied in the small intestine of normal and red bean kidney lectin-treated suckling piglets. In the duodenum and jejunum (but not ileum) of lectin-treated animals overexpression of CB was noted in chromogranin A-immunoreactive (CgA-IR) neuroendocrine (NE) cells. In both control and experimental group a small population of CB-IR NE cells exhibited the presence of somatostatin (but not serotonin, histamine or CRF). After the lectin treatment, an increased (however not statistically significant) immunoreactivity to CB was found in a small subpopulation of neurons of outer submucous (but not inner submucous and myenteric) plexus. It is suggested that there is a functional interaction between lectin administration and CB-expression in the porcine small intestine. Future studies will be needed to clarify this processes.

**Key words:** calbindin, neuroendocrine cells, enteric nervous system, red kidney bean lectin, small intestine, pig

## Introduction

Calbindin 28kDa (CB) belongs to the large EF-hand family of cytosolic proteins which, because of their high affinities for calcium, are called calcium-binding proteins (CaBP). Using molecular techniques structure of the mammalian (including human) cDNAs and genes coding for CB have been identified,

analyzed and cloned (Lomri et al. 1989, Varghese et al. 1989, Parmentier 1990). Several studies have revealed the wide distribution of CB in mammalian tissues, predominantly in central and peripheral nervous system. The expression of CB gene was detected in discrete regions of the rat brain including cells of dentate gyrus, inferior olivary nucleus, the nuclei of the trapezoid body, habenular nuclei, entorhinal cortex,

the mammillary nuclei as well as in forebrain, brainstem gray matter and the posterior horn of the spinal cord (Abe et al. 1992). Different amounts of CB-immunoreactive (CB-IR) neurons and nerve fibres were additionally noted in the mammalian sympathetic (Masliukov et al. 2012), parasympathetic (Kuramoto et al. 1990, Richardson et al. 2003), sensory (Kuramoto et al. 1990) and enteric ganglia (Furness et al. 1990, Houghton et al. 1992, Arciszewski et al. 2009). However, neurons are not the only one source of CaBP since the expression of CB was also determined in many types of non-neuronal cells like ependymal cells (Abe et al. 1992), renal collecting tubule cells and the cortical collecting duct cells (Pollock and Santiesteban 1995), osteoblastic cells (Bellido et al. 2000), enterochromafin-like (ECL) cells and other subsets of neuroendocrine cells (Pochet et al. 1989, Katsetos et al. 1994, Andersson et al. 1998, Timurkaan and Tarakci 2007). It is generally thought that CaBP act as “trigger” or “buffers”. CB is believed to belong to the latter group and as a regulator of the intracellular  $Ca^{2+}$  level is involved in variety of pivotal processes (for review see Schmidt 2012).

Weaning is one of the most stressful and critical period in lifetime of every mammalian species (with a special emphasis on the pig). Due to biochemical complicity of the weaning process, diet change, remodeling of the gut structure (maturation) and physiology as well as because of the decreased immunity leading to viral infections in the post-weaned piglets anorexia and in consequence relatively high mortality is noted (Madec et al. 2000, Lallès 2008, Grau-Roma et al. 2012). High economic losses in pork production are the main reason why many bio-scientists are involved in searching for an effective way to improve the ratio of successfully weaned pigs. In last years, red bean kidney (*Phaseolus vulgaris*) lectin has been added to the narrow list of useful commercial plant extracts which given orally prior to the weaning hasten the gut maturation and evoke beneficial changes in the gut physiology (Linderoth et al. 2006, Valverde Piedra et al. 2006). Although some basic aspects of lectin activities, like changes in villi morphology (Linderoth et al. 2006) and intestinal absorption (Rådberg et al. 2001), increase in the number of mucosal immune CD19+ and CD3+ cells (Prykhod'ko et al. 2009) or shift in chemical content of enteric neurons (Zacharko-Siembida et al. 2013) have been recently revealed many others are still obscure. Therefore, in the present study we applied immunohistochemical methods to evaluate the changes in the expression patterns of CB-like immunoreactivity in porcine small intestine occurring after the lectin treatment. In the co-localization studies the presence of serotonin, histamine, corticotrophin-releasing factor (CRF), and

somatostatin (SOM) in CB-immunoreactive (IR) neuroendocrine NE cells were investigated in normal and lectin-treated piglets.

## Materials and Methods

*Animals and tissue preparation.* All experimental procedures and animal care received approval from the Local Ethics Committee (20/2012) and stayed in agree with Principles of Laboratory Animal Care, NIH publications No. 86-23, revised 1985. All efforts were made to minimize the number of animals used and their suffering. Ten crossbreed Polish Landrace × Pietrain piglets (weighing approx. 4 kg) of both sexes, born at term and housed in standard farming conditions were used in the study. Three-day-old piglets were divided into a control (n=5) and an experimental group (n=5). In the experimental piglets oral dose of 160 HU/kg b.w. of a red kidney bean lectin preparation according to Pusztai and Watt (1974) was given. Control pigs were given 1 ml of saline to mimic the volume of the lectin administration. After 10 days the control and experimental piglets were sedated with intramuscular injection of azaperone (Stresnil, Janssen-Cilag GmbH, Germany; 0.5 mg/kg b.w.) and transported to the laboratory and killed by an overdose of sodium pentobarbital (Morbital, Biowet Pulawy, Poland; 50 mg/kg b.w.). Portions (approx. 10 cm) from all segments of the small intestine (duodenum, jejunum and ileum) were dissected out and rinsed in cold (4°C) 0.01 M phosphate-buffered saline (PBS; pH=7.3). Tissue samples were opened longitudinally, stretched, pinned onto a piece of balsa wood with serosal surface up and fixed by immersion for 48 h in cold (4°C) Stefanini's solution (containing paraformaldehyde and saturated picric acid). After the fixation the samples were placed for several days in cold 10% sucrose-containing cryoprotective Tyrode's solution (1 change per day). Finally the materials were mounted on a wooden block, embedded in O.C.T. compound and left to freeze in dry ice. Using a cryostat, serial longitudinal and transverse sections of 10 µm thickness were made. Every fifth section was placed on a glass slide (SuperFrost Plus, Menzel GmbH & CoKG, Germany) and stored at -20°C for further immunohistochemical studies.

*Immunohistochemistry.* An indirect immunohistochemical procedure described elsewhere was applied (Arciszewski and Zacharko 2004). Briefly, the slides were dried out at room temperature (30 min, RT) and rehydrated in PBS. Next, the sections were immersed in a blocking solution of PBS containing 10% normal goat serum, 0.25% bovine serum albumin and 0.25% Triton X-100 (Sigma-Aldrich, MO, USA)

at RT. The sections were then incubated overnight in a humid chamber (RT) with a mixture of antibodies raised in two different species. In order to visualize calbindin 28 kDa (CB) in enteric neurons, mouse antibodies raised against general neuronal marker Hu C/D proteins (1:800; Molecular Probes, OR, USA; code A-21271) and rabbit anti-CB antibodies (1:2500; SWant, Switzerland; code CB 38) were applied. The presence of CB in NE cells was confirmed with the use of mouse anti-CB sera (1:2500; SWant, Switzerland; code 300) and rabbit antibodies raised against, widely expressed in subsets of NE cells, chromogranin A (CgA; 1:100, Abcam, UK; code ab45179). In the co-localization studies rabbit (or mouse if necessary) anti-CB antibodies were also mixed with mouse anti-serotonin (1:100; Abcam, UK, code 16007), rabbit anti-corticotrophin releasing factor (CRF; 1:80 Sigma-Aldrich, Germany, code C5348), rabbit anti-histamine (1:600; Pierce Thermo Scientific, USA; code PA118036) or rat anti-somatostatin (SOM; 1:300, Biogenesis, UK, code 8330-0009) antibodies. Next, after three further washes in PBS, the sections were then incubated (1 hour, RT) with a mixture of appropriate secondary antibodies. Texas Red-conjugated anti-rabbit goat IgG (1:400; MP Biomedicals, OH, USA), FITC-conjugated anti-rat goat IgG (1:400; MP Biomedicals) as well as FITC-conjugated anti-mouse goat IgG (dilution 1:400; MP Biomedicals) served as secondary antisera. The slides were finally rinsed in PBS (3x15 minutes) and covered with buffered glycerol (pH 7.4) supplemented with 0.1% para-phenylenediamine (Sigma-Aldrich, St. Louis, MO, USA) as an anti-fading factor. The observations were accomplished by using a spinning disk confocal microscope (BX-DSU Olympus, Nagano, Japan) equipped with appropriate interference filters for detection of Texas Red (545-580 nm; MWIY2) and FITC (470-490 nm; MNIBA2). Expression of two fluorescent levels was studied by interchanging filters. All images were captured using a digital color camera (DP-70, Olympus) and Cell<sup>M</sup> software (Olympus). The specificity of primary antibodies was demonstrated by the preabsorption experiments, in which control sections were incubated with primary antibodies that had been inactivated with an excess of synthetic corresponding antigens (10-100 µg of blocking substance per 1 ml of diluted antiserum). The omission of primary antibodies, their substitution by normal, non-immune sera were also applied as a negative control. In all stained control sections no specific immunoreactions was observed.


*Quantification and statistical analysis.* In all portions of the small intestine of control and experimental animals, the numbers of CB-expressing NE cells

were assessed by cell counting according to the following protocol. In each section stained with CB, lamina muscularis mucosae was first identified and measured. The integrity of lamina muscularis mucosae and mucosa was also checked. Then, in the sections with no damaged and interrupted mucosa (during sectioning) and lamina muscularis mucosae of the length not shorter than 10 mm, all CB-IR NE cells were identified and counted. In each segment of the small intestine, the mean from at least 5 slides from each animal (n=5 in control and n=5 in experimental group) was calculated. The numbers of CB-IR NE cells found in 1 cm of mucosa were expressed as a mean ± SEM. The visual semi-quantitative evaluation of the numbers of CB-IR NE cells was additionally done according to the following five-grade scale: absent, single, moderate, numerous and very numerous. The proportions of CB-IR NE cells co-expressing the biologically active substances (serotonin or SOM) were determined by cell counting and presented as a percentage relative to the total number of CB-IR NE cells. In each animal a random sample of no less than two hundreds of CB-IR NE cells was counted. The mean from at least 5 slides from each animal (n=5 in control and n=5 in experimental group) was calculated. The numbers of CB-IR enteric neurons were expressed as a percentage relative to the total number of Hu C/D expressing neurons. The obtained data were presented as mean ± SEM. The distribution patterns and densities of nerve fibres expressing CB were assessed visually according to the scale presented above. The one-way analysis of variance test (ANOVA) followed by Bonferroni's post hoc test were used to compare data between control and experimental groups. Differences were considered as significant at  $P < 0.05$ .

## Results

*General description.* In the duodenum, jejunum and ileum from the control and experimental animals, the expression of CB was found in different numbers of NE cells. CB-IR NE cells were intensively stained, easily distinguishable from the background, and presented very characteristic morphology (Fig. 1A, 2C, 2D). The majority of CB-IR NE cells possessed axon-like extended cytoplasmic processes, however oval cells with no or very short and blunt extensions were also seen. In animals from the control and experimental groups, all CB-IR NE cells were immunopositive to CgA (Fig. 1A). In control and experimental animals, the presence of CB was also found in certain proportions of enteric neurons and nerve fibres of the small intestine (Fig. 1D).







*Expression of CB and co-localization of CB with serotonin, histamine, CRF, SOM in NE cells.* In the duodenal mucosa of the control piglets, CB-IR NE cells were numerous ( $529 \pm 44$ ,  $n=5$ ; Fig. 1B). When compared to the control, the number of CB-IR NE cells in the mucosa of lectin-treated animals ( $907 \pm 85$ ,  $n=5$ ; Fig. 1C) was statistically higher ( $P<0.05$ ). In both control and experimental piglets, the vast majority of CB-IR NE cells were found in the middle portion of the intestinal villi. CB-IR NE cells located in the duodenal crypts as well as in apical portion of the duodenal villi were moderately numerous. In the jejunal mucosa of the control piglets, CB-IR NE cells were predominantly seen in the middle portion of the villi (Fig. 2A). Lectin administration statistically increased ( $P<0.05$ ) the average number of CB-IR NE cells present in 1 cm of the jejunal mucosa (to  $263 \pm 44$  vs.  $148 \pm 32$  found in control group;  $n=5$ ; Fig. 2B) but had no influence on the distribution of CB-IR NE cells throughout villi. In sections from the jejunum of both control and experimental piglets, CB-IR NE cells were very rare in crypts and apical portion of villi. In control pigs, the mucosa of the ileum contained single CB-IR NE cells exclusively located in the middle portions of intestinal villi. After the lectin treatment the number of CB-IR NE cells present in the ileal mucosa was statistically unchanged when compared to the control ( $57 \pm 19$  vs.  $42 \pm 22$ , respectively). In the ileal mucosa of the lectin-treated animals no CB-IR NE cells located in crypts as well as apical portion of villi were found. In all segments of the small intestine of both experimental as well as control piglets, none of CB-IR NE cells exhibited the presence of serotonin (Fig. 2C) as well as CRF (Fig. 2D). Small subpopulations of CB-IR NE cells additionally co-storing SOM were found in the duodenum, jejunum and ileum of control animals ( $4.6 \pm 1.1\%$ ,  $3.9 \pm 0.8\%$  and  $5.1 \pm 0.9\%$  respectively;  $n=5$ ). When compared to the control, statistically similar ( $P<0.05$ ) proportions of CB-IR/SOM-IR NE cells were noted in the duodenum ( $3.7 \pm 1.0\%$ ;  $n=5$ ), jejunum ( $4.4 \pm 0.8\%$ ;  $n=5$ ) and ileum ( $4.1 \pm 1.2\%$ ;  $n=5$ ) of the lectin-treated piglets (Fig. 2E). In control as well as lectin-treated piglets immunoreactivity to histamine has not been detected in NE cells, including those being CB-IR.

*CB in enteric nervous system.* In both control and experimental group, different proportions of CB-ex-

pressing neurons were detected in myenteric plexus (MP) as well as outer submucous plexus (OSP). In animals from both groups none of neurons located in inner submucous plexus (ISP) exhibited the presence of CB (Fig. 1D). In experimental and control piglets, moderate numbers of CB-IR nerve fibres were found to run between neurons of MP and OSP (but not ISP). In the duodenum, jejunum and ileum of the control animals as many as  $14.6 \pm 3.4\%$ ,  $18.3 \pm 3.2\%$  and  $19.4 \pm 4.0\%$  (respectively) Hu-IR/CB-IR myenteric neurons were found. No statistical changes were noted in percentages of CB-IR myenteric neurons in lectin-treated piglets ( $16.6 \pm 3.2\%$  in duodenum,  $18.8 \pm 3.7\%$  in jejunum and  $20.4 \pm 2.9\%$  in ileum). When compared to the control animals increased proportion of CB-IR OSP neurons were found in the duodenum ( $8.4 \pm 3.4\%$  vs.  $4.9 \pm 1.2\%$  in control) and jejunum ( $8.0 \pm 1.3\%$  vs.  $4.7 \pm 0.6\%$  in control), however these changes appeared to be statistically irrelevant. In both control and experimental animals the proportions of CB-IR OSP neurons found in the ileum were statistically similar ( $6.7 \pm 1.1\%$  and  $6.9 \pm 1.2\%$  respectively). After the lectin treatment no visible changes in density and distribution patterns of CB-IR nerve fibres were noted.

## Discussion

In the present study the abundant and region-specific expression of CB was observed in the small intestine of the normal piglets. IR to CB was limited to enteric neurons/nerve fibres and NE cells, however the numbers of CB-IR NE cells decreased gradually in aboral direction. The lectin stimulation substantially increased the numbers of CB-IR NE cells in the duodenum and jejunum (but not in the ileum). For several reasons CB-IR NE cells observed in the control and lectin-treated small intestine closely resembles ECL cells of the stomach. ECL cells are defined as NE cells of morphology similar to enterochromaffin (EC) cells, located exclusively in the oxintic part of the stomach, storing and secreting histamine (but not serotonin) and expressing CgA. Since virtually all ECL cells also co-store CB, several authors proposed to use this CaBP as a specific marker of the gastric ECL cells (Furness et al. 1989, Andersson



Fig. 1. Expression of CB in the small intestine of the normal and lectin-treated piglets. (A) In double IHC stainings of the duodenum of the lectin-treated piglets the presence of CgA (red colour) in CB-IR neuroendocrine cells (green colour) is marked with arrows (CB-IR/CgA-IR cells are seen as yellow because of red and green colour fusion). The increased numbers of duodenal CB-IR NE cells (arrows) after the lectin treatment is illustrated in (B) and (C). In lectin-treated piglets the increased expression of CB was also found in certain subpopulation of duodenal and jejunal enteric neurons (D); CB-IR neurons are marked with arrows, MP – myenteric plexus, OSP – outer submucous plexus; ISP – inner submucous plexus.




Fig. 2. CB-immunoreactivity in NE cells of the small intestine of the lectin-treated piglets. When compared to the control (A), in the jejunum of the lectin-treated piglets the increased proportion of CB-IR NE cells (arrows) was seen (B). Illustrations of the co-localization studies are presented in (C), (D) and (E). In the jejunum neither serotonin (C) nor CRF (D) was present in CB-IR NE cells. In (E) three different subpopulations of the duodenal NE cells are marked with: arrows (CB-IR/SOM-IR), arrowheads (CB-positive/SOM-negative) and empty arrowheads (CB-negative/SOM-positive).

et al. 1998). Taking into account the above it must be pointed out that, surprisingly, CB-IR NE cells identified in the present study fulfill some of criteria given for ECL cells (except location and histamine-expression). In the present study we found numerous CgA-IR/CB-IR NE cells with a shape similar to EC cells but since none of CB-IR NE cells expressed serotonin they definitely can not be considered as EC cells. Significance of this finding is unknown yet, however it may suggest the presence in the porcine small intestine a highly specialized subpopulation of NE cells. Further immunohistochemical analysis revealed that both in control and lectin-stimulated pigs nearly the same small subpopulation of CB-IR NE cells co-express SOM (but not CRF) which indicates that at least two different subpopulations of CB-IR NE cells can be distinguished. Noteworthy, in ECL cells of the rat, guinea-pig, dog and cat no CB/SOM co-localization was found (Buffa et al. 1989). In the small intestine, SOM secreted from D-cells is recognized as inhibitory neurohormone down-regulating the secretion of other hormones and playing a role in suppression of the gut contractility and blood flow circulation (see Van Op den Bosch et al. 2009 for review). In patients with inflammatory bowel disease the numbers of SOM-IR NE cells is substantially decreased (Watanabe et al. 1992). Therefore, it is possible that in CB-IR/SOM-IR NE cells SOM may inhibit CB release. In fetal sheep, the co-existence of CB and SOM, gastrin and glucagon (but not glucose dependent insulinotropic peptide) in duodenal NE cells was observed (Houghton et al. 1992). Peripheral CRF secreted also from NE cells has been recognized as a modulator of intestinal disturbances occurring during the stress (Taché and Bonaz 2007) which may suggest that increased proportion of CB-positive/CRF-negative NE cells observed in the lectin-treated pigs is not stress-related.

It is widely known that one of the most important function of NE cells is recognition of luminal nutrients and appropriate secretory response in order to enhance food absorption but also to adapt to unfavorable conditions (Buchan 1999). Therefore, the numbers and chemical content of NE cells can be plastic during some pathological processes like inflammation or parasite infection (Lomax et al. 2006). In the present study we demonstrated that lectin stimulation also alters the number of CB-IR NE cells. Although the precise explanation of this finding has to be elucidated such situation from some points of view seems to be beneficial for the neonates development. Since in the small intestine CB was found to facilitate calcium transport across the mucosa (Wassermann and Taylor 1966) the higher calcium uptake would hypothetically explain a better skeleton calcification occurring after

the lectin-treatment (Simmons et al. 1975, Izbicka et al. 1997, Kartsogiannis et al. 2008). During the weaning an increased apoptosis in epithelial cells is noted which frequently leads to disintegration of the intestinal barrier and viral infections (Zabielski et al. 2008). In this light, one of the key problem during weaning is to reduce apoptosis ratio. Apoptotic cell death-preventing properties of CB have been reported in relation to several cell types including PC12 cells (McMahon et al. 1998), osteocytes and osteoblasts (Liu et al. 2004), pancreatic  $\beta$ -cells (Rabinovitch et al. 2001) and neurons as well (Sun et al. 2011). Whether CB has a role in protecting epithelial cells from apoptotic cell death has not been reported yet, but higher expression of CB in NE cells after the lectin treatment may support this hypothesis. Neuroprotective properties of CB and its role against the excitotoxicity were widely reported in relation to many types of neurons (Rintoul et al. 2001, Yenari et al. 2001, Kim et al. 2010, Yuan et al. 2013). Noteworthy, in the present study we, after the lectin treatment, also found an increase (however not statistically significant) of CB content in certain subpopulation of submucous (but not myenteric) neurons what may suggest that lectin, at least to some degree, mediates CB-dependent promotion of survival of enteric neurons. In previous report, an increased levels of galanin and nitric oxide synthase (but not vasoactive intestinal polypeptide and substance P) were noted in the small intestine submucous neurons of the lectin-treated piglets (Zacharko-Siembida et al. 2013).

In conclusion, we herein described for the first time the changes in CB expression in the small intestine of the lectin-treated piglets. After the lectin stimulation increased numbers of CB-IR/CgA-IR cells were noted in the duodenum and jejunum (but not in the ileum). In co-localization studies the existence of SOM (but not serotonin and CRF) was revealed in a subpopulation of CB-IR NE cells. An increase of CB in a small supopulation of duodenal and jejunal OSP neurons was also determined. Our results may provide a new clue on the mechanisms underlying the weaning process, piglets welfare and maintenance. It is suggested that in the porcine small intestine, lectin stimulating CB-expression may be an effective and beneficial factor influencing various physiological aspects of GIT adaptation before weaning.

### Acknowledgements

The excellent technical assistance in cryosectioning of Mr. Roman Lalak is gratefully acknowledged.



## References

- Abe H, Watanabe M, Yamakuni T, Kuwano R, Takahashi Y, Kondo H (1992) Localization of gene expression of calbindin in the brain of adult rats. *Neurosci Lett* 138: 211-215.
- Andersson J, Mei Q, Sundler F (1998) Immunohistochemical markers for enterochromaffin-like (ECL) cells of the rat stomach. *Biomed Res* 19: 227-236.
- Arciszewski MB, Zacharko A (2004) Origin and chemical coding of primary afferent neurones supplying the prostate of the dog. *Anat Histol Embryol* 33: 326-333.
- Arciszewski MB, Całka J, Wasowicz K, Majewski M (2009) Distribution and chemical coding of calretinin- and calbindin-expressing enteric neurons in the duodenum of the sheep. *Pol J Vet Sci* 12: 423-431.
- Bellido T, Huening M, Raval-Pandya M, Manolagas SC, Christakos S (2000) Calbindin-D28k is expressed in osteoblastic cells and suppresses their apoptosis by inhibiting caspase-3 activity. *J Biol Chem* 275: 26328-26332.
- Buchan AM (1999) Nutrient Tasting and Signaling Mechanisms in the Gut III. Endocrine cell recognition of luminal nutrients. *Am J Physiol* 277: G1103-G1107.
- Buffa R, Mare P, Salvatore M, Solcia E, Furness JB, Lawson DE (1989) Calbindin 28 kDa in endocrine cells of known or putative calcium-regulating function. Thyro-parathyroid C cells, gastric ECL cells, intestinal secretin and enteroglucagon cells, pancreatic glucagon, insulin and PP cells, adrenal medullary NA cells and some pituitary (TSH?) cells. *Histochemistry* 91: 107-113.
- Furness JB, Padbury RT, Baimbridge KG, Skinner JM, Lawson DE (1989) Calbindin immunoreactivity is a characteristic of enterochromaffin-like cells (ECL cells) of the human stomach. *Histochemistry* 92: 449-451.
- Furness JB, Trussell DC, Pompolo S, Bornstein JC, Smith TK (1990) Calbindin neurons of the guinea-pig small intestine: quantitative analysis of their numbers and projections. *Cell Tissue Res* 260: 261-272.
- Grau-Roma L, Stockmarr A, Kristensen CS, Enøe C, López-Soria S, Nofrartas M, Bille-Hansen V, Hjulsgaard CK, Sibila M, Jorsal SE, Fraile L, Baekbo P, Vigre H, Segalés J, Larsen LE (2012) Infectious risk factors for individual postweaning multisystemic wasting syndrome (PMWS) development in pigs from affected farms in Spain and Denmark. *Res Vet Sci* 93: 1231-1240.
- Houghton PE, Buchan AM, Challis JR (1992) Ontogeny of the distribution and colocalization of calbindin D28K within neural and endocrine cells of the gastrointestinal tract of fetal and neonatal sheep. *Regul Pept* 37: 73-83.
- Izbicka E, Dunstan CR, Horn D, Adams R, Mundy GR (1997) Mitogenic lectin concanavalin A induces calvarial bone formation in vivo via indomethacin-sensitive pathway. *Calcif Tissue Int* 60: 204-209.
- Kartsogiannis V, Sims NA, Quinn JM, Ly C, Cipetic M, Poulton IJ, Walker EC, Saleh H, McGregor NE, Wallace ME, Smyth MJ, Martin TJ, Zhou H, Ng KW, Gillespie MT (2008) Osteoclast inhibitory lectin, an immune cell product that is required for normal bone physiology in vivo. *J Biol Chem* 283: 30850-30860.
- Katsetos CD, Jami MM, Krishna L, Jackson R, Patchefsky AS, Cooper HS (1994) Novel immunohistochemical localization of 28,000 molecular-weight (Mr) calcium binding protein (calbindin-D28k) in enterochromaffin cells of the human appendix and neuroendocrine tumors (carcinoids and small-cell carcinomas) of the midgut and foregut. *Arch Pathol Lab Med* 118: 633-639.
- Kim SA, Jeon JH, Son MJ, Cha J, Chun MH, Kim IB (2010) Changes in transcript and protein levels of calbindin D28k, calretinin and parvalbumin, and numbers of neuronal populations expressing these proteins in an ischemia model of rat retina. *Anat Cell Biol* 43: 218-229.
- Kuramoto H, Furness JB, Gibbins IL (1990) Calbindin immunoreactivity in sensory and autonomic ganglia in the guinea pig. *Neurosci Lett* 115: 68-73.
- Lallès JP (2008) Nutrition and gut health of the young pig around weaning: what news? *Arch Zootech* 11: 5-15.
- Linderth A, Prykhod'ko O, Ahrén B, F4k F, Pierzynowski SG, Weström BR (2006) Binding and the effect of the red kidney bean lectin, phytohaemagglutinin, in the gastrointestinal tract of suckling rats. *Br J Nutr* 95: 105-115.
- Liu Y, Porta A, Peng X, Gengaro K, Cunningham EB, Li H, Dominguez LA, Bellido T, Christakos S (2004) Prevention of glucocorticoid-induced apoptosis in osteocytes and osteoblasts by calbindin-D28k. *J Bone Miner Res* 19: 479-490.
- Lomax AE, Linden DR, Mawe GM, Sharkey KA (2006) Effects of gastrointestinal inflammation on enteroendocrine cells and enteric neural reflex circuits. *Auton Neurosci* 126-127: 250-257.
- Lomri N, Perret C, Gouhier N, Thomasset M (1989) Cloning and analysis of calbindin-D28K cDNA and its expression in the central nervous system. *Gene* 80: 87-98.
- Madec F, Eveno E, Morvan P, Hamon L, Blanchard P, Cariolet R, Amenna N, Morvan H, Truong C, Mahe D, Albina E, Jestin A (2000) Post-weaning multisystemic wasting syndrome (PMWS) in pigs in France: clinical observations from follow-up studies on affected farms. *Liv Prod Sci* 63: 223-233.
- Masliukov PM, Korobkin AA, Nozdrachev AD, Timmermans JP (2012) Calbindin-D28k immunoreactivity in sympathetic ganglionic neurons during development. *Auton Neurosci* 167: 27-33.
- McMahon A, Wong BS, Iacopino AM, Ng MC, Chi S, German DC (1998) Calbindin-D28k buffers intracellular calcium and promotes resistance to degeneration in PC12 cells. *Brain Res Mol Brain Res* 54: 56-63.
- Parmentier M (1990) Structure of the human cDNAs and genes coding for calbindin D28K and calretinin. *Adv Exp Med Biol* 269: 27-34.
- Pochet R, Blachier F, Malaisse W, Parmentier M, Pasteels B, Pohl V, Résibois A, Rogers J, Roman A (1989) Calbindin-D28 in mammalian brain, retina, and endocrine pancreas: immunohistochemical comparison with calretinin. *Adv Exp Med Biol* 255: 435-443.
- Pollock AS, Santiesteban HL (1995) Calbindin expression in renal tubular epithelial cells. Altered sodium phosphate co-transport in association with cytoskeletal rearrangement. *J Biol Chem* 270: 16291-16301.
- Prykhod'ko O, Fed'kiv O, Linderth A, Pierzynowski SG, Weström BR (2009) Precocious gut maturation and immune cell expansion by single dose feeding the lectin phytohaemagglutinin to suckling rats. *Br J Nutr* 101: 735-742.
- Pusztai A, Watt WB (1974) Isolectins of *Phaseolus vulgaris*. A comprehensive study of fractionation. *Biochim Biophys Acta* 365: 57-71.
- Rabinovitch A, Suarez-Pinzon WL, Sooy K, Strynadka K, Christakos S (2001) Expression of calbindin-D(28k) in a pancreatic islet beta-cell line protects against

- cytokine-induced apoptosis and necrosis. *Endocrinology* 142: 3649-3655.
- Rådberg K, Biernat M, Linderöth A, Zabielski R, Pierzynowski SG, Westrom BR (2001) Enteral exposure to crude red kidney bean lectin induces maturation of the gut in suckling pigs. *J Anim Sci* 79: 2669-2678.
- Richardson RJ, Grkovic I, Anderson CR (2003) Immunohistochemical analysis of intracardiac ganglia of the rat heart. *Cell Tissue Res* 314: 337-350.
- Rintoul GL, Raymond LA, Baimbridge KG (2001) Calcium buffering and protection from excitotoxic cell death by exogenous calbindin-D28k in HEK 293 cells. *Cell Calcium* 29: 277-287.
- Schmidt H (2012) Three functional facets of calbindin D-28k. *Front Mol Neurosci* 5: 25.
- Simmons DJ, Lesker PA, Ellsasser JC (1975) Survival of osteocompetent marrow cells in vitro and the effect of PHA-stimulation on osteoinduction in composite bone grafts. *Proc Soc Exp Biol Med* 148: 986-990.
- Sun S, Li F, Gao X, Zhu Y, Chen J, Zhu X, Yuan H, Gao D (2011) Calbindin-D28K inhibits apoptosis in dopaminergic neurons by activation of the PI3-kinase-Akt signaling pathway. *Neuroscience* 199: 359-367.
- Taché Y, Bonaz B (2007) Corticotropin-releasing factor receptors and stress-related alterations of gut motor function. *J Clin Invest* 117: 33-40.
- Timurkaan S, Tarakci G (2007) Distribution of calbindin-D28k in endocrine cells of gastrointestinal tract and pancreas and some endocrine glands (thyroid, parathyroid, pituitary, adrenals) in rats. *Revue Med Vet* 158: 229-233.
- Valverde Piedra JL, Kapica M, Szymanczyk-Kwapik SE, Pawlowska M, Puzio I (2006) The effect of oral lectin administration on the structure of the gastrointestinal mucosa of newborn rats. *Folia Univ Agric Stetin Zootech* 48: 81-86.
- Van Op den Bosch J, Adriaensen D, Van Nassauw L, Timmermans JP (2009) The role(s) of somatostatin, structurally related peptides and somatostatin receptors in the gastrointestinal tract: a review. *Regul Pept* 156: 1-8.
- Varghese S, Deaven LL, Huang YC, Gill RK, Iacopino AM, Christakos S (1989) Transcriptional regulation and chromosomal assignment of the mammalian calbindin-D28k gene. *Mol Endocrinol* 3: 495-502.
- Wasserman RH, Taylor AN (1966) Vitamin d3-induced calcium-binding protein in chick intestinal mucosa. *Science* 152: 791-793.
- Watanabe T, Kubota Y, Sawada T, Muto T (1992) Distribution and quantification of somatostatin in inflammatory disease. *Dis Colon Rectum* 35: 488-494.
- Yenari MA, Minami M, Sun GH, Meier TJ, Kunis DM, McLaughlin JR, Ho DY, Sapolsky RM, Steinberg GK (2001) Calbindin d28k overexpression protects striatal neurons from transient focal cerebral ischemia. *Stroke* 32: 1028-1035.
- Yuan HH, Chen RJ, Zhu YH, Peng CL, Zhu XR (2013) The neuroprotective effect of overexpression of calbindin-D(28k) in an animal model of Parkinson's disease. *Mol Neurobiol* 47: 117-122.
- Zabielski R, Godlewski MM, Guilloteau P (2008) Control of development of gastrointestinal system in neonates. *J Physiol Pharmacol* 59 (Suppl 1): 35-54.
- Zacharko-Siembida A, Valverde Piedra JL, Szymańczyk S, Arciszewski MB (2013) Immunolocalization of NOS, VIP, galanin and SP in the small intestine of suckling pigs treated with red kidney bean (*Phaseolus vulgaris*) lectin. *Acta Histochem* 115: 219-225.