

# ITS sequence data and morphology differentiate *Cytospora chrysosperma* associated with trunk disease of grapevine in northern Iran

Mahdi Arzanlou\*, Abolfazl Narmani

Plant Protection Department, Faculty of Agriculture, University of Tabriz, P.O. Box 5166614766, Tabriz, Iran

Received: October 10, 2014

Accepted: March 16, 2015

**Abstract:** Trunk diseases are potential threats for the grapevine industry owing to the worldwide incidence and economic impact of the diseases. Several fungal groups are known to be involved in these diseases. In a survey on grapevine trunk diseases in northern Iran, *Cytospora* isolates were repeatedly recovered from vines showing decline symptoms. The symptoms appeared as pale brown to brown streaks in longitudinal cuts of shoots. The morphological and cultural characteristics of the isolates were in agreement with the description of *Cytospora chrysosperma*. Sequence data of the ITS-rDNA region was used to further confirm the identity of the species. Phylogenetic analysis based on the sequence data obtained in this study and the sequences from GenBank, confirmed the morphological identification. Our isolates were clustered together with *C. chrysosperma* isolates known from other woody host plant species. The pathogenicity assay on detached shoots of grapevines induced the same symptoms as was observed in field conditions. Although, *C. chrysosperma* is known from several woody hosts in Iran, the occurrence of this species on grapevines showing decline symptoms is new. The economic impact, distribution, and degree of involvement of *C. chrysosperma* in decline of vines in other regions of Iran remains to be studied.

**Key words:** brown streaks, decline, ITS-rDNA, perennial canker, *Valsa sordaria*

## Introduction

The Iranian grapevine industry plays a significant role in the country's economy. According to the statistics of the United Nations Food and Agriculture Organization (FAO), the Iranian grapevine industry presently covers a total area of 286,000 ha, with an annual production of 3,000,000 t (FAO 2012). The Iranian viticulture system is mainly traditional. The establishment of new vineyards is done differently compared to other countries. In Iran, vineyards are routinely established by direct planting of rooted grapevine cuttings. The grafting process is not a common practice.

Grapevines are susceptible to a large variety of devastating diseases caused by fungi, bacteria, viruses, and other biotic and abiotic agents. Trunk diseases constrict grapevine cultivation and vineyard productivity and longevity in many countries (Mugnai *et al.* 1999; Mostert *et al.* 2006; Arzanlou *et al.* 2013a). The economic importance of trunk diseases on the grapevine industry led to the establishment of an international council on grapevine trunk disease (IGTD). The aim of IGTD is to promote collaboration and the exchange of information among scientists involved in research on grapevine trunk diseases. A biannual workshop is organized by the IGTD on grapevine trunk diseases. The 10th workshop on grapevine trunk diseases was held in November 2014, in Australia. The

term fungal trunk diseases, is attributed to the group of diseases whose agents are located in the mature stem (trunk and cordons) of grapevines. Esca and related diseases caused by *Phaeoacremonium* and *Phaeoconiella*, represent historic and well-studied trunk diseases on grapevines (Mugnai *et al.* 1999; Mostert *et al.* 2006; Essakhi *et al.* 2008; Arzanlou *et al.* 2013a; Arzanlou *et al.* 2014). Other diseases are caused by such fungal pathogens as *Eutypa*, *Botryosphaeria*, *Cylindrocarpon*, and *Phomopsis* (Phillips 1998, 2002; Luque *et al.* 2005, 2006; Van Niekerk *et al.* 2004, 2005, 2011; Taylor *et al.* 2005; Alves *et al.* 2008; Trouillas and Gubler 2010; Arzanlou *et al.* 2012). Grapevine trunk diseases are generally considered to be complex diseases in which a combination of fungal groups are involved (Mostert *et al.* 2006). A brief overview on the etiologic agents of grapevine trunk diseases is presented in table 1. In recent years, several other fungal groups have been characterised from grapevines with trunk disease symptoms, including: *Cadophora luteo-olivacea*, *Macrophomina phaseolina*, *Pestalotiopsis uvicola*, *Verticillium dahliae*, *Phomopsis viticola*, *Truncatella angustata*, *Fusarium* spp., *Cytospora* spp., and *Acremonium* spp. However, the contribution and degree of involvement for many of the above-mentioned fungal groups in trunk-disease progress on grapevines remain unknown and their pathogenicity on grapevines still need to be tested (Trouillas and Gubler

\*Corresponding address:  
Arzanlou@tabrizu.ac.ir

**Table 1.** Fungal species known as the etiologic agents of grapevine trunk diseases

Black foot	Eutypa dieback	Esca/young vine decline	Bot canker
<i>Campylocarpon fasciculare</i>	<i>Eutypa leptoplaca</i>	<i>Phaeoacremonium aloesii</i>	<i>Diplodia corticola</i>
<i>Campylocarpon pseudofasciculare</i>	<i>Eutypa laevata</i>	<i>Phaeoacremonium angustius</i>	<i>Diplodia mutila</i>
<i>Cylindrocarpon pauciseptatum</i>	<i>Eutypella cryptovalsoidea</i>	<i>Phaeoacremonium armeniacum</i>	<i>Diplodia seriata</i>
<i>Cylindrocarpon liriodendra</i>	<i>Eutypella microtheca</i>	<i>Phaeoacremonium australiense</i>	<i>Dothiorella</i> sp.
<i>Cylindrocarpon macrodidymum</i>	<i>Eutypa leptoplaca</i>	<i>Phaeoacremonium austroafricanum</i> = <i>Truneatella austroafricana</i>	<i>Dothiorella americana</i>
<i>Cylindrocarpon obtusisporum</i>	<i>Eutypella vitis</i>	<i>Phaeoacremonium cinereum</i>	<i>Lasiodiplodia crassispora</i>
<i>Ilyonectria macrodidyma</i>	<i>Eutypella</i> spp.	<i>Phaeoacremonium croatiense</i>	<i>Lasiodiplodia missouriana</i>
<i>Ilyonectria liriodendra</i>	<i>Cryptosphaeria lygniota</i>	<i>Phaeoacremonium globosum</i>	<i>Lasiodiplodia theobromae</i>
<i>Ilyonectria alcacerensis</i>	<i>Cryptosphaeria pullmanensis</i>	<i>Phaeoacremonium griseorubrum</i>	<i>Lasiodiplodia viticola</i>
<i>Ilyonectria estremocensis</i>	<i>Cryptovalsa ampelina</i>	<i>Phaeoacremonium hispanicum</i>	<i>Neofusicoccum australe</i>
<i>Ilyonectria novozelandica</i>	<i>Cryptovalsa rabenhorstii</i>	<i>Phaeoacremonium hungaricum</i>	<i>Neofusicoccum luteum</i>
<i>Ilyonectria torresensis</i>	<i>Diatrype brunneospora</i>	<i>Phaeoacremonium inflatipes</i>	<i>Neofusicoccum macroclavatum</i>
	<i>Diatrype oregonensis</i>	<i>Phaeoacremonium iraniamum</i>	<i>Neofusicoccum mediterraneum</i>
	<i>Diatrype stigma</i>	<i>Phaeoacremonium krajdenui</i> = <i>T. krajdenui</i>	<i>Neofusicoccum parvum</i>
	<i>Diatrype whitmanensis</i>	<i>Phaeoacremonium mortoniae</i> = <i>T. fraxinopennsylvanica</i>	<i>Neofusicoccum ribis</i>
	<i>Diatrypella verrucaeformis</i>	<i>Phaeoacremonium occidentale</i>	<i>Neofusicoccum viticlavatum</i>
	<i>Diatrypella vulgaris</i>	<i>Phaeoacremonium parasiticum</i> = <i>T. parasitica</i>	<i>Neofusicoccum vitifusiforme</i>
		<i>Phaeoacremonium rubrigenum</i> = <i>T. rubrigena</i>	<i>Phaeobotryosphaeria porosa</i>
		<i>Phaeoacremonium sicilianum</i>	<i>Spencermartinsia viticola</i>
		<i>Phaeoacremonium scolyti</i>	<i>Neofusicoccum ribis</i>
		<i>Phaeoacremonium subulatum</i>	
		<i>Phaeoacremonium tuscanum</i>	
		<i>Phaeoacremonium venezuelense</i>	
		<i>Phaeoacremonium viticola</i> = <i>T. viticola</i>	

2010; Arzanlou *et al.* 2012; Arzanlou *et al.* 2013b; Arzanlou *et al.* 2014).

Species of the genus *Cytospora* are generally known as the causal agents of canker (referred to as Valsa canker, Leucostoma canker or perennial canker) and dieback on woody host plant species, with rare occurrences on herbaceous plants. Over 85 woody hosts have been listed as susceptible to *Cytospora* canker including many economically important fruit trees as well as shade and ornamental tree species in different stages of growth (Sinclair *et al.* 1987; Fotouhifar *et al.* 2007, 2010). *Cytospora* canker is considered to be a destructive disease on stone fruit trees especially on peach (Barakat and Johnson 1997). The pathogens infect the inner bark (secondary phloem) and the outer bark (periderm) (Adams *et al.* 2005, 2006). Some of the *Cytospora* spp. are known to have an endophytic life style, colonising xylem tissues in healthy woody hosts (González and Tello 2011).

A number of *Cytospora* spp. are known to have a wide host range, occurring on a diverse range of plant species e.g. *C. chrysosperma* and *C. cincta*; while others are restricted to members of one plant family or a genus e.g. *C. sacchari* and *C. tritici* (Adams *et al.* 2005, 2006). There is a lack


of knowledge on the occurrence and contribution of *Cytospora* species in grapevine trunk diseases worldwide. González *et al.* (2011) have listed *C. chrysosperma* as an endophyte colonising grapevine woody tissues in Spain.

Several fungal species have been reported to occur on grapevines in Iran and have been reviewed by Arzanlou *et al.* (2013a). The first aim of this study was to characterise *Cytospora* spp. associated with grapevine canker disease in northwestern Iran by means of morphological and molecular data. The second aim was to further evaluate the pathogenicity of the agents using a detached shoot assay.

## Materials and Methods

### Isolates and morphology


Wood samples were collected from grapevines with decline symptoms. The symptoms included stunted growth, leaf yellowing and necrosis, internal wood necrosis, black vascular streaking or discoloured tissues (Figs. 1 and 2). The samples were collected in vineyards of the northwestern zone of Iran including the East and West Azerbaijan provinces. Isolation was made following the proto-



**Fig. 1.** External and internal decline symptoms on grapevines from which only *Cytospora chrysosperma* was recovered: a–d – stunted growth and leaf necrosis in a vineyard; e–l – various types of wood necrosis and discoloration in cross sections through the wood

col described by Arzanlou and Dokhanchi (2013a, b), and Arzanlou *et al.* (2012, 2013a, b, 2014). In brief, small pieces, approximately  $0.5 \times 0.5 \times 0.5$  cm, were cut from the discoloured wood tissues and surface-sterilised for 15–20 sec in 70% ethanol, rinsed three times with sterile water, dried on sterile filter paper, and transferred to Potato Dextrose Agar (PDA, Fluka, Hamburg, Germany) plates supplemented with  $100 \text{ mg} \cdot \text{l}^{-1}$  streptomycin sulphate and  $100 \text{ mg} \cdot \text{l}^{-1}$  ampicillin. Pure cultures were established

by employing the hyphal tip technique. The cultures were preserved on PDA in 2 ml microtube slants at  $4^\circ\text{C}$  in the Culture Collection of Tabriz University (CCTU). The morphological characteristics were examined based on pure cultures. Colony morphology, including colour, shape, and growth rate, was determined after 7 days of incubation on PDA at  $25^\circ\text{C}$  in darkness. Microscopic characters were studied using a smash mount technique with sterile distilled water, as explained by Arzanlou *et al.*



**Fig. 2.** External and internal decline symptoms on grapevines from which together with *Cytospora chrysosperma* other fungal trunk pathogens were also recovered: a–c – stunted growth and leaf necrosis in a vineyard; d–h – various types of wood necrosis and discoloration in cross sections through the wood

(2007). The dimensions of microscopic structures were calculated based on 30 measurements for conidia morphology (shape, colour, and cell number) and size (length and width) where possible.

### Molecular identification

#### DNA extraction

For DNA extraction, fungal isolates were grown on PDA for 8 days in the dark. Fresh mycelia were collected and subjected to DNA extraction by using the protocol of Moller *et al.* (1992).

#### Sequence analysis

The ITS (International Transcribed Spacer) region, including the 3' end of the 18S rRNA gene, ITS1, 5.8S rDNA, ITS2, and the 5' end of 28S rRNA gene, was amplified using the primer set V9G (Vilgalys and Hester 1990) and ITS4 (White *et al.* 1990). The reaction mixture was the same as for that of Arzanlou and Dokhanchi (2013a, b), Arzanlou and Khodaei (2012). The reaction was performed on a GeneAmp PCR System 9700 (Applied Biosystems, Foster City, CA) with cycling conditions consisting of 5 min at 96°C for primary denaturation, followed by 40 cycles of denaturation at 94°C for 30 sec, annealing at 52°C for 30 sec, extension at 72°C for 60 sec, with a final extension at 72°C for 7 min. Sequencing of the Polymerase Chain Reaction (PCR) products was done using BigDye Terminator v3.1 (Applied Biosystems, Foster City, CA) Cycle Sequencing Kit according to the manufacturer's recommendation. The analysis was done on an ABI Prism 3700 (Applied Biosystems, Foster City, CA). Raw sequence files were edited manually by using SeqMan™II (DNASTAR, Madison, Wisconsin, USA), and a consensus sequence was generated for each of the sequences. Sequences were subjected to a Megablast search analysis at NCBI's GenBank nucle-

otide database for sequence similarity. Sequences with a high similarity were downloaded from GenBank. The sequences were aligned by using ClustalW algorithm implemented in MEGA 5 (Tamura *et al.* 2011). The automatically aligned file was further checked by eye. A minimum number of gaps were inserted to improve the alignment. Phylogenetic trees were constructed using neighbour-joining method with the following criteria: substitution model as Jukes-Cantor; gap treatment as pairwise deletion and transitions and transversions (with the equal ratio). The supports of the internal nodes of the trees were evaluated by the bootstrap method with 10,000 replicates. *Diaporthe CBS 267* (GenBank accession number: KC343081.1) was used as outgroup in the phylogenetic analyses.

#### Pathogenicity studies

Excised shoot assay was used to evaluate pathogenicity of the isolates in laboratory conditions as described by Arzanlou *et al.* (2012). In brief, 1-year old shoots, 10–15 mm in diameter and 25 cm in length, were cut from a healthy mature vein of cv. Askari. The leaves were removed and shoots were surface-sterilised with 70% ethanol. For inoculation, shoots were wounded 10 cm above the first internode by removing the cortex with a sterile 5 mm diameter metal cork borer. A mycelial plug (5 mm diameter) obtained from the margin of a seven-day-old fungal colony was placed in the wound with the mycelium facing the stem. The wound was wrapped with Parafilm and for the control shoot a plug of MEA was placed in the wound. Inoculated shoots were placed in a plastic container containing moistened filter papers to keep the relative humidity high. The shoots in the containers were kept in laboratory conditions at 25°C with natural daylight. Shoots were examined after 28 days for disease progress. Longitude sections were made going upward and downward from the inoculation point with *C. chrysosperma* isolate. The length of necrosis was used as an indicator of pathogenicity. The

experiment was carried out using four fungal isolates and three replicates for each treatment.

## Results

### Disease symptoms

The symptoms on grapevines appeared as typical decline symptoms. The symptoms included stunted growth, die-back and leaf chlorosis, and necrosis. Internal symptoms appeared as different degrees of wood discolouration in the transverse section though brown wood discoloura-

tion in sapwood was evident. From the symptoms presented in figure 1, only *C. chrysosperma* was recovered. From the symptoms shown in figure 2, besides *C. chrysosperma*, other fungal trunk pathogens were also isolated.


### DNA phylogeny and morphology

Phylogeny inferred using the sequence data of the ITS-rDNA region from the isolates obtained in this study (Table 2), with other *Cytospora* spp. from GenBank, clustered our isolates with *C. chrysosperma* (100% bootstrap support value) (Fig. 3).

**Table 2.** The list of *Cytospora chrysosperma* isolates used for phylogenetic analysis in this study

Collection number	Country of origin
<sup>a</sup> CCTU153	<sup>b</sup> WA, Iran
CCTU154	WA, Iran
CCTU155	WA, Iran
CCTU156	WA, Iran
CCTU383	WA, Iran
CCTU400	WA, Iran
CCTU401	WA, Iran
CCTU402	WA, Iran
CCTU405	WA, Iran
CCTU406	WA, Iran

<sup>a</sup> Culture Collection of Tabriz University, Iran; <sup>b</sup> West Azerbaijan province




**Fig. 3.** A neighbour-joining phylogenetic tree obtained from the ITS region and 5.8S rDNA sequence data. Bootstrap support values (> 65) from 1,000 replicates are indicated on the nodes. The tree was rooted to *Diaporthe eres* (GenBank number: KC343081.1). The scale bar indicates 0.05 substitutions per site

The cultural and morphological characteristics of our isolates were in full agreement with the description for *C. chrysosperma* (Adams *et al.* 2006). Colonies attained a diameter of 40 mm after 7 days of incubation at 25 on PDA. Colonies were flat to slightly raised, felty, and with entire margins. Colony colour on the surface was a moderate yellowish and on the reverse the colour was a moderate to strong yellow to moderate orange yellow. Conidiomata produced abundantly on PDA after two weeks of incubation. Conidiomata were scattered, labyrinthine cytosporoid, with labyrinthine chambers, circular to ovoid, 1–2 mm in diameter. Locules were multi-chambered, and often irregular. Conidiophores were hyaline, and of two types. Type 1 were branched and septate, 15–30  $\mu\text{m}$  in length. Type 2 were long, unbranched, and aseptate or rarely septate. Conidiogenous cells were phialidic, subcylindric, and hyaline. Conidia were hyaline, allantoid, and aseptate, 3.5–5(4)  $\times$  1  $\mu\text{m}$ , oozing out as an orange mucilaginous mass (Fig. 4).


#### Pathogenicity studies

The results of the pathogenicity assay revealed that *C. chrysosperma* is pathogenic on detached shoots of

grapevines. The shoots were examined after 28 days to note the progress of the disease. Longitudinal and cross sections were made from above and below the inoculation point. Wood necrosis was evident on the inoculated shoots. No obvious symptoms were observed in the control. After removing the bark, a superficial brown discoloration was observed up and down from the inoculation points. In cross sections through the inoculation points, there was some wood discoloration apparent in some cases. There were substantial differences among the isolates in the length of lesions induced on shoots. The isolates D and C induced the largest lesion areas on detached shoots with a mean value of  $12 \pm 2.64$  and  $12.33 \pm 1.2$  cm, respectively; while, the lengths of the lesion induced by isolates A and B were  $6.50 \pm 1.04$  and  $7.16 \pm 1.01$  cm, respectively. Pycnidia were developed on inoculated shoots. No fungal structure was observed on the control (Fig. 5). Koch's postulates were fulfilled by the re-isolation of the causal agents from the inoculated shoots. *Cytospora chrysosperma* was recovered only from the inoculated shoots. No fungal growth was observed in the controls (Fig. 5).



**Fig. 4.** *Cytospora chrysosperma*: a – colony morphology on PDA; b–c – cross section through conidiomata on PDA; d – conidiomata on bark of inoculated shoot; e–f – cross section through conidiomata developed on artificially inoculated shoot; g – conidiophores; h – conidia. Scale bars: c, e = 200  $\mu\text{m}$ ; f = 100  $\mu\text{m}$ ; g, h = 10  $\mu\text{m}$



**Fig. 5.** Pathogenicity assay using excised shoot method: a–b – pycnidium on inoculated shoot; c – superficial wood necrosis upward and downward from the inoculation point; d–g – disease symptoms and discoloration on inoculated shoot with *Cytospora chrysosperma*; h – control with no obvious disease symptom

## Discussion

Trunk diseases are among the most destructive diseases of vineyards worldwide. Several fungal groups have been noted on grapevines with trunk diseases symptoms (Table 1). In the present study, *Cytospora* isolates were isolated from grapevines displaying typical symptoms of decline including stunted growth, leaf chloro-

sis and necrosis. The other fungal groups isolated from symptomatic tissues (Fig. 2) included *Togninia minima* (asexual morph: *Phaeocremonium aleophilum*) (Arzanlou and Narmani 2014), *Fusarium* spp., *Acremonium* spp., and Pestalotid fungi. Various types of internal wood discoloration were observed in cross sections through the cane and trunk. Morphological and molecular char-

acterization of *Cytospora* isolates revealed the identity of species as *C. chrysosperma*. *Cytospora chrysosperma* is known to have a wide host range with common occurrence on members of Salicaceae, and is rare on other woody angiosperms throughout the world (Gvritshvili 1982; Hayova and Minter 1998). Fotouhifar *et al.* (2007, 2010) have provided a list of the host plant species for *C. chrysosperma* in Iran. There are only a few reports on the occurrence of *Cytospora* species on grapevines. Fotouhifar *et al.* (2010) have reported *C. cincta* and *C. leucostoma* from grapevine in Iran. González and Tello (2011) have listed *C. chrysosperma* as an endophyte colonising grapevine woody tissues in Spain.

The inoculation studies carried out in this study, showed *C. chrysosperma* isolates being pathogenic on the excised shoot of grapevine. The results on the pathogenicity of *Cytospora* spp. are controversial. On *Prunus* (Biggs 1989) and *Populus* species (Kepley and Jacobi 2000), *Cytospora* species have proven to be highly virulent and destructive. However, in general, *Cytospora* species are considered as wound parasites attacking trees weakened by biotic and abiotic factors (Schoeneweiss 1975, 1983). Some of the species have an endophytic lifestyle colonising bark and xylem tissues of woody hosts (Fisher *et al.* 1993; Bills 1996).

After putting all of the results together, it can be said that *C. chrysosperma* is a potential pathogen on grapevine in Iran. Further studies on the host range, pathogenicity, and virulence of this species on different cultivars of *Vitis vinifera* are now required.

## Acknowledgements

The authors would like to thank the Research Deputy of the University of Tabriz and the Studienstiftung Mykologische Systematik und Ökologie for their financial support.

## References

- Adams G.C., Roux J., Wingfield M.J. 2006. *Cytospora* species (Ascomycota, Diaporthales, Valsaceae): introduced and native pathogens of trees in South Africa. *Australasian Plant Pathology* 35 (5): 521–548.
- Adams G.C., Wingfield M.J., Common R., Roux J. 2005. Phylogenetic relationships and morphology of *Cytospora* species and related teleomorphs (Ascomycota, Diaporthales, Valsaceae) from *Eucalyptus*. *Studies in Mycology* 52: 1–142.
- Alves A., Crous P.W., Correia A., Phillips A.J.L. 2008. Morphological and molecular data reveal cryptic speciation in *Lasiodiplodia theobromae*. *Fungal Diversity* 28 (1): 1–13.
- Arzanlou M., Dokhanchi H. 2013a. Phenotypic and molecular characterization of *Diplodia seriata*, the causal agent of canker and twig dieback disease on mulberry in Iran. *Archives of Phytopathology and Plant Protection* 46 (6): 682–694.
- Arzanlou M., Dokhanchi H. 2013b. Calosphaeria canker of almond caused by *Calosphaeria pulchella* in Iran. *Archives of Phytopathology and Plant Protection* 46 (2): 215–226.
- Arzanlou M., Khodaei S. 2012. *Aureobasidium iranianum*, a new species on bamboo from Iran. *Mycosphere* 3 (4): 404–408.
- Arzanlou M., Moshari S., Bakhshi M., Khodaei S. 2012. First report of *Botryosphaeria dothidea* associated with grapevine decline disease in Iran. *Australasian Plant Disease Notes* 7 (1): 197–200.
- Arzanlou M., Moshari S., Salari M., Badali H. 2013a. Molecular characterization and pathogenicity of *Phaeoacremonium* spp. associated with esca disease of grapevine in Northern Iran. *Archives of Phytopathology and Plant Protection* 46 (4): 375–388.
- Arzanlou M., Narmani A., Moshari S., Khodaei S., Bababi-ahari A. 2013b. *Truncatella angustata* associated with grapevine trunk disease in Northern Iran. *Archives of Phytopathology and Plant Protection* 46 (10): 1168–1181.
- Arzanlou M., Narmani A. 2014. Multiplex PCR assay for specific identification and determination of mating type applied to *Togninia minima* (anamorph known as *Phaeoacremonium aleophilum*), one of the causal agents of esca disease in grapevine. *Phytopathology Mediterranean* 53 (2): 240–249.
- Arzanlou M., Narmani A., Moshari S., Khodaei S. 2014. Pome and stone fruit trees as possible reservoir hosts for *Phaeoacremonium* spp., the causal agents of grapevine esca disease, in Iran. *Archives of Phytopathology and Plant Protection* 47 (6): 717–727.
- Barakat R.M., Johnson D.A. 1997. Expansion of cankers caused by *Leucostoma cincta* on sweet cherry trees. *Plant Disease* 81 (12): 1391–1394.
- Biggs A.R. 1989. Integrated approach to control *Leucostoma* canker of peach in Ontario. *Plant Disease* 73 (11): 869–874.
- Bills G.F. 1996. Isolation and analysis of endophytic fungal communities from woody plants. p. 31–65. In: "Endophytic Fungi in Grasses and Woody Plants: Systematics, Ecology, and Evolution" (S.C. Redlin, L.M. Carris, eds.). APS Press, St. Paul, MN, USA, 321 pp.
- Essakhi S., Mugnai L., Crous P.W., Groenewald J.Z., Surico G. 2008. Molecular and phenotypic characterization of novel *Phaeoacremonium* species isolated from esca diseased grapevines. *Persoonia* 21: 119–134.
- FAO (Food and Agriculture Organization of the United Nations). 2012. FAO Statistical Databases. Available at <http://www.fao.org/>
- Fisher P.J., Petrini O., Sutton B.C. 1993. A comparative study of fungal endophytes from leaves, xylem and bark of *Eucalyptus nitens* in Australia and England. *Sydowia* 45 (2): 338–345.
- Fotouhifar K.B., Hedjaroude G.A., Ershad D., Moussavi S.M., Okhovvat S.M., Javan Nikkhah M. 2007. New information on the form-genus *Cytospora* in Iran (I). *Rostaniha* 8 (2): 129–149.
- Fotouhifar K.B., Hedjaroude G.A., Leuchtmann A. 2010. ITS rDNA phylogeny of Iranian strains of *Cytospora* and associated teleomorphs. *Mycologia* 102 (6): 1369–1382.
- Gvritshvili M.N. 1982. The fungal genus *Cytospora* in the USSR. *Izdatelstve Sabchota Sakarstvelo*, Tbilisi, Russia, 214 pp.
- González V., Tello M.L. 2011. The endophytic mycota associated with *Vitis vinifera* in central Spain. *Fungal Diversity* 47 (1): 29–42.
- Hayova V.P., Minter D.W. 1998. *Leucostoma cinctum*. *IMI Descriptions of Fungi and Bacteria*. Set 137 No. 1361, 3 pp.
- Kepley J.B., Jacobi W.R. 2000. Pathogenicity of *Cytospora* fungi on six hardwood species. *Journal of Arboriculture* 26 (6): 326–332.
- Luque J., Martos S., Phillips A.J.L. 2005. *Botryosphaeria viticola* sp. nov. on grapevines: A new species with a *Dothiorella* anamorph. *Mycologia* 97 (5): 1111–1121.



- Luque J.D., Sierra E., Torres E., Garcia F. 2006. *Cryptovalsa ampelina* on grapevines in N.E. Spain: identification and pathogenicity. *Phytopathology Mediterranea* 45 (Suppl.): 101–109.
- Moller E.M., Bahnweg G., Geiger H.H. 1992. A simple and efficient protocol for isolation of high molecular weight DNA from filamentous fungi, fruit bodies, and infected plant tissues. *Nucleic Acids Results* 20 (22): 6115–6116.
- Mostert L., Groenewald J.Z., Summerbell R.C., Gams W., Crous P.W. 2006. Taxonomy and pathology of *Togninia* (Diaporthales) and its *Phaeoacremonium* anamorphs. *Studies of Mycology* 54: 1–115.
- Mugnai L., Graniti A., Surico G. 1999. Esca (black measles) and brown wood streaking: two old and elusive diseases of grapevines. *Plant Disease* 83 (5): 404–418.
- Phillips A.J.L. 1998. *Botryosphaeria dothidea* and other fungi associated with excoriose and dieback of grapevines in Portugal. *Journal of Phytopathology* 146: 327–332.
- Phillips A.J.L. 2002. *Botryosphaeria* species associated with diseases of grapevines in Portugal. *Phytopathology Mediterranea* 41 (1): 3–18.
- Schoeneweiss D.F. 1975. Predisposition, stress and plant disease. *Annual Review of Phytopathology* 13: 193–211.
- Schoeneweiss D.F. 1983. Drought predisposition to *Cytospora* canker in blue spruce. *Plant Disease* 67: 383–385.
- Sinclair W.A., Lyon H.H., Johnson W.T. 1987. *Diseases of Trees and Shrubs*. Cornell University Press Ithaca, NY, USA, 680 pp.
- Tamura K., Nei M., Kumar S. 2011. MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 28 (10): 2731–2739.
- Taylor A., Hardy G.E.S., Wood P., Burgess T. 2005. Identification and pathogenicity of *Botryosphaeria* species associated with grapevine decline in Western Australia. *Australasian Plant Pathology* 34 (2): 187–195.
- Trouillas F.P., Gubler W.D. 2010. Pathogenicity of Diatrypaceae species in grapevines in California. *Plant Disease* 94 (7): 867–872.
- Van Niekerk J.M., Bester W., Halleen F., Crous P.W., Fourie P.H. 2011. The distribution and symptomatology of grapevine trunk disease pathogens are influenced by climate. *Phytopathology Mediterranea* 50: S98–S111.
- Van Niekerk J.M., Crous P.W., Groenewald J.Z., Fourie P.H., Halleen F. 2004. DNA phylogeny, morphology and pathogenicity of *Botryosphaeria* species on grapevines. *Mycologia* 96 (4): 781–798.
- Van Niekerk J.M., Groenewald J.Z., Farr D.F., Fourie P.H., Halleen F., Crous P.W. 2005. Reassessment of *Phomopsis* species on grapevines. *Australasian Plant Pathology* 34 (1): 27–39.
- Vilgalys R., Hester M. 1990. Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172 (8): 4238–4246.
- White T.J., Bruns T., Lee S., Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. p. 315–322. In: "PCR Protocols: a Guide to Methods and Applications" (M. Innis, D. Gelfand, J. Sninsky, T. White, eds.). Academic Press, Orlando, Florida, 482 pp.