

ISOLATION AND CHARACTERIZATION OF AN *ARABIDOPSIS THALIANA* SELF-INCOMPATIBILITY MUTANT INDUCED BY HEAVY-ION BEAM IRRADIATION

KOK SONG LAI^{1*} AND TAKEHISA MASATSUGU²

¹*Agro-Biotechnology Institute Malaysia, Ministry of Science, Technology and Innovation, c/o MARDI Headquarters, 43400 Serdang, Selangor DE, Malaysia*

²*Graduate School of Biological Sciences, Nara Institute of Science and Technology, 8916-5 Takayama, Ikoma 630-0192, Japan*

Received December 17, 2012; revision accepted November 21, 2013

Self-incompatibility (SI) is a genetic system that promotes outcrossing by rejecting self-pollen. In the Brassicaceae the SI response is mediated by the pistil S-locus receptor kinase (SRK) and its ligand, pollen S-locus cysteine-rich (SCR) protein. Transfer of *SRK-SCR* gene pairs to self-fertile *Arabidopsis thaliana* enabled establishment of robust SI, making this transgenic self-incompatible *A. thaliana* an excellent platform for SI analysis. Here we report isolation of a novel *A. thaliana* self-incompatibility mutant, AtC24 SI mutant, induced by heavy-ion beam irradiation. We show that the AtC24 SI mutant exhibits breakdown of SI, with pollen hydration, pollen tube growth and seed set resembling the corresponding processes in wild-type (self-fertile) *A. thaliana*. Further reciprocal crosses indicated that some perturbed SI factor in the stigmatic cell of the AtC24 SI mutant is responsible for the observed phenotype, while the pollen response remained intact. Our results demonstrate successful application of heavy-ion beam irradiation to induce a novel *A. thaliana* self-incompatibility mutant useful for SI studies.

Key words: *Arabidopsis thaliana* accession C24, heavy-ion beam, pollen, pollen tube, self-incompatibility.

INTRODUCTION

In many angiosperms, outcrossing is enforced by genetic self-incompatibility (SI), which allows the pistil to recognize and specifically reject self-pollen (Tantikanjana and Nasrallah, 2012). In the Brassicaceae (crucifer) family, the SI reaction is based on allele-specific interactions between two highly polymorphic proteins encoded by the S-locus haplotype: S-locus receptor kinase (SRK) displayed on the surface of stigma epidermal cells (Takasaki et al., 2000), and its ligand S-locus cysteine-rich (SCR) protein localized in the pollen coat (Schopfer et al., 1999; Takayama et al., 2000). Under one current model of SI signalling, contact between a pollen grain and stigma epidermal cell allows binding of SCR to its cognate SRK receptor to take place and activate the downstream SI cascades, leading to rejection of self-pollen.

Recently, the natural self-compatible (self-fertile) model plant *Arabidopsis thaliana* has been

made to express SI upon transformation with *SRKb-SCRb* gene pairs derived from its self-incompatible sister species *A. lyrata* (Nasrallah et al., 2002). This finding implies that *A. thaliana* has retained all the required downstream components of SI signalling, making this self-incompatible *A. thaliana* (hereafter SI *Arabidopsis*) an excellent platform for SI analysis. Several SI molecules have been suggested to function downstream of *Brassica* SI (Stone et al., 1999; Murase et al., 2004; Samuel et al., 2009) but it has also been suggested that *A. thaliana* orthologs of *Brassica* SI molecules are not required for the SI response of *A. thaliana* (Kitashiba et al., 2011). Hence the molecular events precipitated by activation of SRK in *A. thaliana* SI are still unknown. Further studies through mutagenesis and subsequent mapping of the target gene are needed to identify the downstream SI molecules of *A. thaliana*.

Heavy-ion beam irradiation has often been employed as an efficient mutagenic technique to induce a desired mutant for functional gene analy-

*e-mail: christian_lks@yahoo.com




Fig. 1. Schematic map of *SRKb-SCRb/pBI121* expression plasmid. RB – right border; NOS-pro – nopaline synthase promoter; NPT II – neomycin phosphotransferase; NOS-t – nopaline synthase terminator; SCR-pro – S-locus cysteine-rich promoter; SRK-pro – S-locus receptor kinase promoter; LB – left border.

sis. Successful isolation of target mutants induced by heavy-ion beam irradiation has been reported for varieties of ornamental plants such as dahlia (Hamatani et al., 2001), petunia (Miyazaki et al., 2002), torenia (Miyazaki et al., 2006) and verbena Kanaya et al., 2008). This powerful technique has also been used to induce mutation in rice (Hayashi et al., 2007), *Arabidopsis* (Kazama et al., 2008b) and tobacco (Kazama et al., 2008a). Unlike gamma rays and X-rays, heavy-ion beams have high linear energy transfer (LET), which exerts stronger biological effects and can produce more DNA damage, leading to mutation and inactivation of single or multiple genes, inducing stable knockout mutants (Kazama et al., 2007). Since a heavy-ion beam induces mutations with high frequency at a relatively low dose, it produces mutants with less cell damage, no severe growth inhibition, and a broad spectrum of mutations (Kazama et al., 2007).

Here we present a preliminary report on our attempt to induce an *A. thaliana* SI mutant (hereafter AtSI mutant) using heavy-ion beam irradiation. We describe a successful application of heavy-ion beam irradiation for isolation and characterization of a novel AtSI mutant which can serve as a platform for downstream SI gene analysis in *A. thaliana*.

MATERIALS AND METHODS

PLANT MATERIAL

Arabidopsis thaliana accession C24 was used throughout this study. Stable SI *Arabidopsis* was generated by transforming the construct harboring *SRKb-SCRb* gene pairs isolated from *A. lyrata* (Fig. 1) into *A. thaliana* C24 using the floral dip method as previously described (Clough and Bent, 1998). Genomic PCR and segregation analysis were performed to confirm the homozygous SI *Arabidopsis*. Gene-specific primers are listed in Table 1.

PLASMID CONSTRUCTION

The expression vector used for *A. thaliana* C24 transformation was constructed by cloning the *SRKb-SCRb* gene pairs from *A. lyrata* into the

pBI121 plasmid driven by its own native promoter (Fig. 1). The *SRKb* gene fragment was inserted into the *Kpn1* site, and the *SCRb* gene fragment was inserted between *Sma1-Sac1* sites, producing *SRKb-SCRb/pBI121* plasmid (Fig. 1). Gene-specific primers are listed in Table 1.

HEAVY-ION BEAM IRRADIATION

Dry SI *Arabidopsis* seeds (~5000 seeds) were packed with Hybri-Bag Hard (95 µm thickness, Cosmo Bio, USA) to provide a monolayer of seeds for homogenous irradiation. Heavy-ion beam irradiation was performed according to the previously described protocol (Kazama et al., 2008b) at the RIKEN Nishina Center Radioactive-Isotope Beam Factory (RIBF) Facility, Saitama, Japan.

ISOLATION OF AtSI MUTANT

The irradiated SI *Arabidopsis* seeds (~1100 seeds) were surface-sterilized and sown on 0.8% (w/v) agar containing 1/2 Murashige and Skoog (MS) medium (Wako, Japan) supplemented with MS vitamins (Sigma Aldrich, USA) and incubated at 25°C under a 16 h photoperiod. A total 1000 seedlings were grown in Arasystem trays (Betatech, Ghent, Belgium) to generate the M_1 population. To obtain seeds for the M_2 population, M_1 plants were incubated with dry ice for 3 h to induce self-fertilization. A total 255 independent M_2 lines were successfully recovered for AtSI mutant screening. To screen for AtSI mutants, silique length of the M_2

TABLE 1. Primers used in the study

Primer name	Sequence (5'-3')
SCRb-F1	CCCGGGATGAGGAATGCTACTTTCTTC
SCRb-R1	GAGCTCTAGCAAATCTACAGTCGCATA
SRKb-F1	AGGTACCATGAGAGTTGTAGTACCAAACCTG
SRKb-R1	TGGTACCTTACCGAGGGTTCGATGGCCGA
SRKb-RT-F1	AAAGAACAGGGGATTCTACAACCTCAA
SRKb-RT-R1	ACTTCATGTGGTTCGAAACGC
GAPDH-F1	GACCTACTGTCTCAGACTCGAG
GAPDH-R1	TTAGGCCCTTGACATGTGGACGA

lines was measured as previously described (Lai et al., 2012a).

CHARACTERIZATION OF AtC24 SI MUTANT

A pollination assay was performed by hand-pollinating pre-emasculated stigmas with pollen grains. Pollen hydration was evaluated after 30 min pollination on papilla cells with pollen grains. Pollen tube growth was monitored on pollinated stigmas using aniline blue staining as described by Sumie et al. (2001). Stained pistils were observed and photographed with a Zeiss Axioplan fluorescence microscope (Carl Zeiss, Oberkochen, Germany).

GENOMIC PCR ANALYSIS AND DNA SEQUENCING

Genomic DNA was extracted from leaves using the DNeasy Plant Mini kit (Qiagen, USA). Genomic PCR analysis was performed according to the protocol described by Lai et al. (2011b, 2012c). DNA sequencing was performed using an ABI 3100 DNA Sequencer (ABI, USA), following the manufacturer's protocol. Primers are listed in Table 1.

QUANTITATIVE REAL-TIME PCR

Total RNA was extracted from stigmas by the previously described protocol (Lai et al., 2011). The concentration and purity of total RNA was determined by spectrophotometric analysis. Quantitative real-time PCR was performed according to the protocol described by Lai et al. (2012b, 2013). GAPDH was the control. Gene-specific primers are listed in Table 1.

RESULTS AND DISCUSSION

As a first step toward isolating the AtSI mutant we established SI *Arabidopsis* suitable for use in heavy-ion beam irradiation. Previously, transformation of *A. thaliana* C24 with *SRK-SCR* gene pairs was shown to produce stable SI *Arabidopsis* expressing robust SI indistinguishable from the naturally self-incompatible *A. lyrata* (Nasrallah et al., 2002). Hence, the *A. thaliana* C24 genetic background was used in this study. An expression vector harboring the *SRKb-SCRb* gene pairs was constructed (Fig. 1) and transformed into *A. thaliana* C24. Homozygous SI *Arabidopsis* exhibiting stable SI was screened using genomic PCR and confirmed through segregation analysis and pollination assay (data not shown). The resulting SI *Arabidopsis* exhibiting developmentally stable SI was used in subsequent heavy-ion beam irradiation.

SI *Arabidopsis* seeds (~5000 seeds) were heavy-ion beam irradiated at the RIBF Facility,




Fig. 2. Outline of screening for AtSI mutant from heavy-ion beam irradiated mutant population.

Saitama, Japan (Fig. 2). Consistent with previously reported data (Kazama et al., 2008b), the irradiated M_1 seeds achieved 92.1% germination (data not shown). This result was expected, as heavy-ion beams induce mutations with high frequency at a relatively low dose without significantly affecting seed germination and plant survivability. A total 1000 M_1 seedlings were grown to generate the M_2 population (Fig. 2). To induce self-fertilization and obtain M_2 seeds the M_1 lines were incubated with dry ice (CO_2 source). Previously, adding carbon dioxide to the environment has been shown to induce seed set in SI *Brassica* genotypes (O'Neill et al., 1988). Carbon dioxide has been suggested to overcome SI by enhancing pollen activity during germination and tube growth (Dhaliwal et al., 1981), blocking the callose response in the stigmatic papilla cell in expression of SI (O'Neill et al., 1984), and increasing the rate of pollen adhesion (Palloix et al., 1985). Of the 1000 M_1 lines incubated, we successfully recovered and grew 255 individual M_2 lines for use in AtSI mutant screening (Fig. 2).

To screen for AtSI mutants, silique length of the M_2 lines was measured. From the 255 M_2 lines screened, we isolated a mutant line with well-developed siliques (mean length 11.0 mm, vs 12.5 mm in wild type; Fig. 3a). This mutant line, the AtC24 SI mutant, produced silique length similar to that seen




Fig. 3. AtC24 SI mutant silique length and pollen tube growth. (a) Length of AtC24 SI mutant siliques. For each sample, 5 siliques were measured in 3 biological replicates, (b) Aniline blue staining of pistils from AtC24 SI mutant. Arrows point to pollen tubes. Bars = 200 μm.




Fig. 4. Aniline blue staining of pistils from reciprocal crosses between SI *Arabidopsis* and AtC24 SI mutant. Arrows point to pollen tubes. Bars = 200 μm.

in self-fertile WT plants (Fig. 3a). No other obvious vegetative growth defects were observed in the AtC24 SI mutant. Further aniline blue staining revealed pollen tube growth in the AtC24 SI mutant (Fig. 3b). Pollen tubes were also observed growing into the style and ovary of the AtC24 SI mutant (Fig. 3b). In the aniline blue staining process, typically only accepted pollen grains will adhere and remain on the stigmatic papillae. All rejected pollen grains failed to adhere to the stigmatic papillae and as a result were washed away during fixation and processing for microscopic examination in the aniline blue staining protocol. Both silique length

measurements and pollen tube monitoring suggest that a factor essential to the SI response was perturbed in the AtC24 SI mutant.

To determine the nature of this SI factor, reciprocal crosses between SI *Arabidopsis* and AtC24 SI mutants were performed. Pollination of SI *Arabidopsis* pistils expressing SRK (hereafter SRK-pistil) with AtC24 SI mutant pollen grains resulted in no pollen tube growth and short siliques (Fig. 4). In contrast, pollination of AtC24 SI mutant pistils with SI *Arabidopsis* pollen grains expressing SCR (hereafter SCR-pollen) resulted in pollen tube growth and well developed siliques (Fig. 4). This




Fig. 5. Characterization of AtC24 SI mutant. (a) Pollen hydration assays of AtC24 SI mutant. For each sample, 25 pollen grains were tested, (b) Seed count of AtC24 SI mutant. For each sample, 5 siliques were counted in 3 biological replicates, (c) Seed set analysis of AtC24 SI mutant. Bars = 1 mm.

result indicates that the SI factor of the female organ (stigma) is compromised in the AtC24 SI mutant. To further determine the extent of pollen acceptance in the AtC24 SI mutant, pollen hydration assay and seed set analysis were performed to assess the degree of SI breakdown. Approximately 96% of the SCR-pollen grains pollinated deposited on AtC24 SI mutant papillae showed successful pollen hydration and germination in the pollen hydration assay (Fig. 5a). This result indicates strong pollen acceptance in the AtC24 SI mutant, further suggesting that the perturbed SI factor is crucial to this SI response. The loss of this SI factor resulted in total breakdown of SI observed in

the AtC24 SI mutant. Similar observations were reported when SI molecules acting downstream of *Brassica* SI signalling were disrupted (Stone et al., 1999; Murase et al., 2004; Samuel et al., 2009). The pollen factor was apparently intact, as no pollen hydration and germination were detected when SRK-pistils were pollinated with AtC24 SI mutant pollen grains (Fig. 5a). In accord with the pollen hydration assay, AtC24 SI mutants with a well developed silique also produced seed set indistinguishable from that of self-fertile WT plants (Fig. 5b,c). No seeds were produced in SI *Arabidopsis* (positive control) (Fig. 5b,c).




Fig. 6. Detection and expression analysis of *SRKb* in AtC24 SI mutant. (a) Genomic PCR of *SRKb* in AtC24 SI mutant. Lane M – 1.0 Kb DNA ladder; lane 1 – *SRKb*-*SCRB*/pBI121 plasmid; lane 2 – SI *Arabidopsis*; lane 3 – AtC24 SI mutant; lane 4 – WT plant, (b) Relative *SRKb* mRNA levels in stigmas of AtC24 SI mutant.

In *Arabidopsis*, heavy-ion beam irradiation has been shown to induce a broad spectrum of mutations, including base changes, small DNA deletions involving a few bases, and large DNA alterations such as inversions, translocations and deletions (Tanaka, 1999). To rule out the possibility that a mutation or deletion in the *SRKb* gene fragment might be responsible for the observed phenotype in the AtC24 SI mutant, genomic PCR analysis and DNA sequencing were performed (Fig. 6). Based on genomic PCR analysis, a ~2.5 Kb fragment coinciding with the size of the *SRKb* gene was detected (Fig. 6a). DNA sequencing did not detect any mutation or deletion in the *SRKb* gene of the AtC24 SI mutant. We also measured the level of the *SRKb* transcript in the AtC24 SI mutant, using qRT-PCR analysis (Fig. 6b). Previously a reduced level of SRK in stigmatic cells has been suggested to cause breakdown of SI (Liu et al., 2007). Quantitative analysis of the *SRKb* transcript level in the AtC24 SI

mutant revealed no significant differences or reduction of *SRKb* expression versus SI *Arabidopsis* (positive control) (Fig. 6b). Thus the breakdown of SI observed in the AtC24 SI mutant was not caused by a mutation or deletion that disrupted the *SRKb* level. Instead, an important novel stigmatic SI factor acting downstream of SRK might be responsible for the breakdown of SI in the AtC24 SI mutant.

Here we reported successful isolation of a novel AtC24 SI mutant induced by heavy-ion beam irradiation. To our knowledge there are no reports on isolation of AtSI mutants induced by heavy-ion beam irradiation. The obtained AtC24 SI mutant will serve as valuable material for SI gene analysis and examination of the downstream SI molecules of *Arabidopsis*.

ACKNOWLEDGMENTS

We thank Dr. Tomoko Abe for irradiating the SI *Arabidopsis* seeds. We are grateful to the members of the Intercellular Communications Laboratory for their constructive suggestions on the study.

REFERENCES

- CLOUGH SJ, and BENT AF. 1999. Floral dip: A simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant Journal* 16: 735–743.
- DHALIWAL AS, MALIK CP, and SINGH MB. 1981. Overcoming incompatibility in *Brassica campestris* L. by carbon dioxide, and dark fixation of the gas by self- and cross-pollinated pistils. *Annals of Botany* 48: 227–233.
- HAMATANI M, IITSUKA Y, ABE T, MIYOSHI K, YAMAMOTO M, and YOSHIDA S. 2001. Mutant flowers of dahlia (*Dahlia pinnata* Cav.) induced by heavy-ion beams. *RIKEN Accelerator Progress Report* 34: 169.
- HAYASHI Y, TAKEHISA H, KAZAMA Y, ICHIDA H, RYUTO H, FUKUNISHI N, ABE T, KAMBA C, and SATO T. 2007. Effects of ion beam irradiation on mutation induction in rice. In: *Eighteenth International Conference, Cyclotrons and Their Applications. Catania, Italy, 1–5 October 2007*, 1–5.
- KANAYA T, SAITO H, HAYASHI Y, FUKUNISHI N, RYUTO H, MIYAZAKI K, KUSUMI T, ABE T, and SUZUKI K. 2008. Heavy-ion beam induced sterile mutants of verbena (*Verbena hybrida*) with an improved flowering habit. *Plant Biotechnology* 25: 91–96.
- KAZAMA Y, SAITO H, FUJIWARA M, MATSUYAMA T, HAYASHI Y, RYUTO H, FUKUNISHI N, and ABE T. 2007. An effective method for detection and analysis of DNA damage induced by heavy-ion beams. *Bioscience Biotechnology, and Biochemistry* 71: 2864–2869.
- KAZAMA Y, SAITO H, MIYAGAI M, TAKEHISA H, ICHIDA H, MIYAZAWA Y, MISHIBA K, KANAYA T, SUZUKI K, BAE CH, MIYOSHI K, MII M, and ABE T. 2008a. Effect of heavy ion-beam irradiation on plant growth mutation induction in *Nicotiana tabacum*. *Plant Biotechnology* 25: 105–111.
- KAZAMA Y, SAITO H, YAMAMOTO YY, HAYASHI Y, ICHIDA H, RYUTO H, FUKUNISHI N, and ABE T. 2008b. LET-dependent effects of

- heavy-ion beam irradiation in *Arabidopsis thaliana*. *Plant Biotechnology* 25: 113–117.
- KITASHIBA H, LIU P, NISHIO T, NASRALLAH JB, and NASRALLAH ME. 2011. Functional test of *Brassica* self-incompatibility modifiers in *Arabidopsis thaliana*. *Proceedings National Academy of Sciences* 108: 18173–18178.
- LAI KS, ABDULLAH P, YUSOFF K, and MAHMOOD M. 2011a. An efficient protocol for particle bombardment-mediated transformation of *Centella asiatica* callus. *Acta Physiologiae Plantarum* 33: 2547–2552.
- LAI KS, KATHOEN-NAKAYAMA P, IWANO M, and TAKAYAMA S. 2012a. A TILLING resource for functional genomics in *Arabidopsis thaliana* accession C24. *Genes and Genetic Systems* 87: 291–297.
- LAI KS, YUSOFF K, and MAHMOOD M. 2011b. Extracellular matrix layer as the early structural marker for *Centella asiatica* embryogenic tissues. *Biologia Plantarum* 55: 549–553.
- LAI KS, YUSOFF K, and MAHMOOD M. 2012b. Functional ectodomain of the hemagglutinin-neuraminidase protein is expressed in transgenic tobacco cells as a candidate vaccine against Newcastle disease virus. *Plant Cell Tissue and Organ Culture* 112: 117–121.
- LAI KS, YUSOFF K, and MAHMOOD M. 2012c. Heterologous expression of hemagglutinin-neuraminidase protein from Newcastle disease virus strain AF2240 in *Centella asiatica*. *Acta Biologica Cracoviensia Series Botanica* 54/1: 1–6.
- LAI KS and YUSOFF K. 2013. Production of tENDO1 in stably transformed tobacco cell culture for mismatch detection. *Plant Cell Tissue and Organ Culture* 114: 287–214.
- LIU P, SHERMAN-BROYLES S, NASRALLAH ME, and NASRALLAH JB. 2007. A cryptic modifier causing transient self-incompatibility in *Arabidopsis thaliana*. *Current Biology* 17: 734–740.
- MIYAZAKI K, SUZUKI K, ABE T, KATSUMOTO Y, YOSHIDA S, and KUSUMI T. 2002. Isolation of variegated mutants of *Petunia hybrida* using heavy-ion beam irradiation. *RIKEN Accelerator Progress Report* 35: 130.
- MIYAZAKI K, SUZUKI K, IWAKI K, ABE T, YOSHIDA S, and FUKUI H. 2006. Flower pigment mutations induced by heavy ion beam irradiation in an interspecific hybrid of *Torenia*. *Plant Biotechnology* 23: 163–167.
- MURASE K, SHIBA H, IWANO M, CHE FS, WATANABE M, ISOGAI A, and TAKAYAMA S. 2004. A membrane-anchored protein kinase involved in *Brassica* self-incompatibility signalling. *Science* 303: 1516–1519.
- NASRALLAH ME, LIU P, and NASRALLAH JB. 2002. Generation of self-incompatible *Arabidopsis thaliana* by transfer of two S locus genes from *A. lyrata*. *Science* 297: 247–249.
- O'NEILL P, SINGH MB, and KNOX RB. 1988. Cell biology of the stigma of *Brassica campestris* in relation to CO₂ effects on self-pollination. *Journal of Cell Science* 89: 541–549.
- O'NEILL P, SINGH MB, NEALES TF, KNOX RB, and WILLIAMS EG. 1984. Carbon dioxide blocks the stigma callose response following incompatible pollinations in *Brassica*. *Plant Cell & Environment* 7: 285–288.
- PALLOIX A, HERVE Y, KNOX RB, and DURNAS C. 1985. Effect of carbon dioxide and relative humidity of self-incompatibility in cauliflower, *Brassica oleracea*. *Theoretical and Applied Genetics* 70: 629–633.
- SAMUEL MA, CHONG YT, HAASEN KE, ALDEA-BRYDGES MG, STONE SL, and GORING DR. 2009. Cellular pathways regulating responses to compatible and self-incompatible pollen in *Brassica* and *Arabidopsis* stigmas intersect at Exo70A1, a putative component of the exocyst complex. *Plant Cell* 21: 2655–2671.
- SCHOPPER CR, NASRALLAH ME, and NASRALLAH JB. 1999. The male determinant of self-incompatibility in *Brassica*. *Science* 286: 1697–1700.
- STONE SL, ARNOLDO M, and GORING DR. 1999. A breakdown of *Brassica* self-incompatibility in ARC1 antisense transgenic plants. *Science* 286: 1736–173.
- SUMIE I, AKIKO K, JUNICHI U, IKUO N, and KIYOTAKA O. 2001. The *DEFECTIVE IN ANTHER DEHISCENCE1* gene encodes a novel phospholipase A1 catalyzing the initial step of jasmonic acid biosynthesis, which synchronizes pollen maturation, anther dehiscence, and flower opening in *Arabidopsis*. *Plant Cell* 10: 2191–2210.
- TAKASAKI T, HATAKEYAMA K, SUZUJI G, WATANABE M, ISOGAI A, and HINATA K. 2000. The S receptor kinase determines self-incompatibility in *Brassica* stigma. *Nature* 403: 913–916.
- TAKAYAMA S, SHIBA H, IWANO M, SHIMOSATO H, CHE FS, KAI N, WATANABE M, SUZUKI G, HINATA K, and ISOGAI A. 2000. The pollen determinant of self-incompatibility in *Brassica campestris*. *Proceedings National Academy of Sciences* 97: 1920–1925.
- TANAKA A. 1999. Mutation induction by ion beams in *Arabidopsis*. *Gamma Field Symposia* 38: 19–27.
- TANTIKANJANA T, and NASRALLAH JB. 2012. Non-cell-autonomous regulation of crucifer self-incompatibility by auxin response factor ARF3. *Proceedings National Academy of Sciences* 109: 19468–19473.